

An aerial photograph of a coastal landscape featuring prominent white cliffs and a sea stack. The cliffs are layered and show signs of erosion. The sea stack is a tall, narrow rock formation in the water. The sky is clear and blue.

A 25 minutes  
overview of

Models of high- $p_T$  parton energy loss  
in a colored medium

Nicolas BORGHINI

CERN TH




Models of high- $p_T$  parton energy loss  
reproduce the data remarkably well







# Models of high- $p_T$ parton energy loss reproduce the data remarkably well








# Models of high- $p_T$ parton energy loss reproduce the data remarkably well






# Models of high- $p_T$ parton energy loss reproduce the data remarkably well






# Models of high- $p_T$ parton energy loss reproduce the data remarkably well







# Models of high- $p_T$ parton energy loss reproduce the data remarkably well





# Models of high- $p_T$ parton energy loss reproduce the data remarkably well





# Models of high- $p_T$ parton energy loss

Welcome to the realm of acronyms!

- ① Radiative vs. collisional **energy loss**
- ① Theories and models of **radiative energy loss**
  - **LPM**-effect based approaches: BDMPS-Z & AMY
  - **opacity** expansion: GLV; (AS)W
  - medium-enhanced **higher-twist** effects
  - medium-modified **MLLA**
- ① Theories and models of **collisional energy loss**






# Models of high- $p_T$ parton energy loss

Two different "categories" of models of parton energy loss, depending on the basic underlying process:

inelastic


"radiative" process (Bremsstrahlung)



also "in vacuum", but controlled by the presence of a medium collisions!

elastic

"collisional" process






# Inelastic energy loss

Models based on the Landau-Pomeranchuk-Migdal effect [1/4]

The propagating **high- $p_T$  parton** traverses a **thick target**.



It radiates **soft gluons**, which scatter **coherently** on **independent color charges** in the medium, resulting in a **medium-modified gluon energy spectrum**.

**Multiple soft scattering limit**



# Inelastic energy loss

Models based on the Landau-Pomeranchuk-Migdal effect [2/4]

Independent scattering centers:  $\lambda \gg 1/\mu$   
 mean free path  $\leftarrow$   $\lambda$   $\gg$   $1/\mu$   $\rightarrow$  screening mass



Note the assumption, which actually underlies all models of in-medium partonic energy loss



Coherent scatterings:  $\frac{l_{\text{coh}}}{k_{\perp}^2} \sim \frac{2\omega}{k_{\perp}^2} \leq L$  (medium length)  
 coherence length  $\leftarrow$   $\frac{l_{\text{coh}}}{k_{\perp}^2}$   $\simeq N_{\text{coh}} \mu^2$   $\Rightarrow$   $l_{\text{coh}} = \sqrt{\frac{2\omega\lambda}{\mu^2}}$   
 of the emitted gluon

LPM only affects gluons with  $\omega \lesssim \omega_c \equiv \frac{1}{2} \hat{q} L^2$

Medium characterized by the transport coefficient  $\hat{q} \equiv \frac{\mu^2}{\lambda}$

Baier, Dokshitzer, Mueller, Peigné, Schiff (BDMPS); Zakharov





# Inelastic energy loss


Models based on the Landau-Pomeranchuk-Migdal effect [3/4]

Gluon coherence length  $l_{\text{coh}} = \sqrt{\frac{2\omega\lambda}{\mu^2}}$

$\Rightarrow$  gluon energy spectrum per unit path length  $\omega \frac{dI}{d\omega dz} \simeq \frac{\alpha_s}{l_{\text{coh}}} \simeq \alpha_s \sqrt{\frac{\hat{q}}{\omega}}$

For a path length  $L$ :  $\omega \frac{dI}{d\omega} \simeq \alpha_s \sqrt{\frac{\hat{q}L^2}{\omega}}$

Average medium-induced energy loss:  $\Delta E = \int^{\omega_c} \omega \frac{dI}{d\omega} d\omega \simeq \alpha_s \omega_c \propto \alpha_s \hat{q} L^2$



👉 BDMPS-Z, only two parameters:  $\hat{q}$  &  $L$



# Inelastic energy loss

Models based on the Landau-Pomeranchuk-Migdal effect [4/4]

What about the infrared ( $\omega \rightarrow 0$ ) behaviour?

👁 BDMPS-Z: **coherent regime** requires

$$N_{\text{coh}} > 1 \Leftrightarrow \ell_{\text{coh}} > \lambda \Leftrightarrow \omega > E_{\text{LPM}} \equiv \lambda \mu^2 = \mathcal{O}(1 \text{ GeV})$$

👁 AMY (Arnold, Moore, Yaffe; Jeon, Gale, Turbide):

interaction of the **fast parton** with a **thermal bath**

✓ **LPM energy loss** for  $\lambda \sim 1/g_s^2 T$ ,  $\mu \sim g_s T \Rightarrow \ell_{\text{coh}} > \lambda \Leftrightarrow \omega \gtrsim T$

✓ and for  $0 < \omega < E_{\text{LPM}} \simeq 1 \text{ GeV}$ , Bethe-Heitler regime

↓  
Energy loss per unit length proportional to the incoming energy

In addition, they allow possible **gains** in the **parton energy**

👉 AMY approach, three **parameters**:  $T$ ,  $L$  &  $\alpha_s$





# Inelastic energy loss

Models based on an **opacity** expansion [1/2]

The **high- $p_T$  parton** interacts with a **thin target**:

the **energy loss** results from an **incoherent superposition** of very few  $\chi \equiv L/\lambda$  single hard scattering processes along the **path length**  $L$ .  
↳ "opacity" (= number of collisions)

⇒ **gluon energy spectrum** per **unit path length**

$$\omega \frac{dI}{d\omega dz} \simeq \left(\frac{L}{\lambda}\right) \frac{\alpha_s}{l_{\text{coh}}} \simeq \left(\frac{L}{\lambda}\right) \alpha_s \frac{\mu^2}{\omega} \neq \alpha_s \sqrt{\frac{\hat{q}}{\omega}} \text{ within LPM}$$

leads to an **average energy loss**  $\Delta E \propto L^2$  (for a **static medium**)

Gyulassy, Lévai, Vitev (GLV); Wiedemann

👉 three parameters:  $\left(\frac{L}{\lambda}\right), \mu$  &  $L$

↳ ⇔ the (linear) **density of scattering centers**





# Inelastic energy loss

Models based on an **opacity** expansion [2/2]

- Within **GLV**, **radiated gluons** restricted to  $\omega > \mu = \mathcal{O}(500 \text{ MeV})$ , “common value” of the **screening mass** and the **plasmon excitation**
- **Energy loss** actually dominated by **energetic gluons**  $\omega \gtrsim \bar{\omega}_c \equiv \frac{1}{2} \mu^2 L$  ( $\neq$  **LPM**, where **soft gluons** with  $\omega < \omega_c$  mainly contribute)
- Only very few ( $\approx 3$ ) **gluons** are radiated by the **fast parton**





# Inelastic energy loss

Approach based on a **twist** expansion

In **QCD**, a cross-section can actually be expanded in powers of  $\frac{1}{q^2}$ , where  $q$  is the exchanged (hard) momentum:

“**twist expansion**”

In vacuum, **higher-twist** terms are power suppressed (!).

But in a **medium**, these terms may become enhanced:  $A^{1/3}/q^2$

⇒ allow systematic computation of energy loss

**formulated** in terms of “**medium**-modified fragmentation functions”  
(which can be evolved with DGLAP...)

Guo, Wang & Wang

Parameters (?):  $\mu, T$





# Inelastic energy loss

A model based on modified parton splitting functions

Effect of the medium modeled by a (phenomenological) modification of the Altarelli-Parisi parton splitting functions, considering e.g.

$$P_{qq}(z) = C_F \left( \frac{2(1 + f_{\text{med}})}{1 - z} - (1 + z) \right)$$

where  $f_{\text{med}} = 0$  in the absence of a medium ( $f_{\text{med}}$  only parameter)

⇒ modification of the “hump-backed plateau” of longitudinal particle distributions within a jet computed using MLLA

NB, Wiedemann

Modified Leading Logarithmic Approximation (of QCD)





# Inelastic energy loss

A model based on **modified parton splitting functions**

Effect of the **medium** modeled by a (phenomenological) modification of the Altarelli-Parisi **parton splitting functions**, considering e.g.


$$P_{qq}(z) = C_F \left( \frac{2(1 + f_{\text{med}})}{1 - z} - (1 + z) \right)$$

where  $f_{\text{med}} = 0$  in the absence of a medium ( $f_{\text{med}}$  only parameter)

⇒ modification of the **“hump-backed plateau”** of longitudinal particle distributions within a **jet** computed using **MLLA**

NB, Wiedemann

depletion  
at large  $x$



enhancement  
at small  $x$




# Inelastic energy loss

A few **model**-independent remarks [1/2]

actually also valid for models of **elastic energy loss**

- 👁 All **partons** do not **lose** the same amount of **energy**, even when they traverse the same **in-medium path length  $L$** 
  - ⇒ **nuclear modification factor  $R_{AA}$**  mostly reflects the few **partons** which have **lost little energy**
  - 👉 use of “**quenching weights**” (= probability to **lose** a given **energy**)
- 👁 The **medium** traversed by the **parton** is not static, but in **expansion!**
  - 👉 **model**-builders introduce **dynamics** (most often, à la Bjorken), which may lead to a redefinition ( $\hat{q} \rightarrow \hat{q}_{\text{eff}}$ ) of the parameters, to the introduction of new ones ( $\tau_0, T_0$ ), or to a change in scaling properties ( $\Delta E_{\text{GLV}} \propto L$  instead of  $L^2$ )





# Inelastic energy loss

A few **model**-independent remarks [2/2]

- A model of **partonic energy loss** has to be supplemented by several **other elements** to allow comparison with the **data**:
  - **parton** distribution functions inside the **nuclei** (**shadowing**, **Cronin effect**...)
  - **production** cross-sections
- ⇒ seemingly similar conclusions of **different models** may actually differ
  - Turbide et al. (**AMY** approach), PRC 72 (2005) 014906:  
reproduce  $R_{AA}$  for pions assuming  $T_i = 370 \text{ MeV}$ ,  $\tau_i = 0.26 \text{ fm}/c$ ,  
 $\frac{dN}{dy} = 1260$  &  $\alpha_s = 0.3$ .  
No need for **initial state effects** as **shadowing** & the **Cronin effect**
  - **GLV**, PRL 89 (2002) 252301:  $\frac{dN^g}{dy} = 1100$   
invoke competition between **shadowing**, **Cronin effect** and **partonic energy loss** to obtain a flat  $R_{AA}$ .






# Inelastic energy loss

Further **model**-dependent remarks [1/3]

Drawing conclusions from **fits** to the **data** may not be easy!

“ $R_{AA}$  is fragile” (Eskola, Honkanen, Salgado, Wiedemann)



Data cannot allow to distinguish between  $\hat{q} = 5$  or  $15 \text{ GeV}^2/\text{fm}$



# Inelastic energy loss

Further **model**-dependent remarks [2/3]

Let me be even more pessimistic / skeptical...

👁 Eskola, Honkanen, Salgado, Wiedemann, NPA 747 (2005) 511:

$$\hat{q} = 5 - 15 \text{ GeV}^2/\text{fm}, \text{ with } \langle L \rangle \simeq 2 \text{ fm}$$

which leads to strong (& questionable?) conclusions

👁 Arleo, hep-ph/0601075:

$$\hat{q} = 0.3 - 0.4 \text{ GeV}^2/\text{fm}, \text{ with } \langle L \rangle \simeq 5 \text{ fm}$$

...but François 1. fixed the latter value a priori & 2. assumed that **all partons lose energy**

👁 Baier & Schiff, hep-ph/0605183:

$$\hat{q} = 1 - 3 \text{ GeV}^2/\text{fm}, \text{ with } \langle L \rangle \simeq 3 \text{ fm}$$

restricting the region of validity of the **LPM effect**





# Inelastic energy loss

Further **model**-dependent remarks [3/3]

Could one compute the **transport coefficient**  $\hat{q}$  ab initio, even in the non-perturbative case?

Idea: use Maldacena's conjecture of a correspondence between **QCD** and its **dual weakly coupled theory of gravity** living in a 5-dimensional anti-de Sitter space-time.

More practically, since the **dual** of **QCD** is unknown, replace it by some **supersymmetric Yang-Mills theory** ("SYM N=4").

$$\hat{q}_{\text{SYM}} = \frac{\pi^2 \sqrt{2} \Gamma(\frac{3}{4})}{\Gamma(\frac{5}{4})} \sqrt{\alpha_{\text{SYM}} N_c} T^3$$

Liu, Rajagopal, Wiedemann

$\hat{q}_{\text{SYM}} \propto \sqrt{N_c} \neq$  number of degrees of freedom is proportional to  $N_c^2$   
 $\rightarrow \Leftrightarrow$  **entropy density**

But... the result is not "**universal**" (may not hold for **QCD**)





# Elastic energy loss

The elder (Bjorken, 1984), yet still in its infancy...

Bjorken (1984), Thoma & Gyulassy (1991), Braaten & Thoma (1991), Wang, Gyulassy & Plumer (1995), Mustafa et al. (1998), Lin, Vogt & Wang (1998):  $dE_{el.}/dz \approx 0.3 - 0.5 \text{ GeV/fm}$ : negligible!

Then, all of a sudden...

Mustafa & Thoma (2003), Dutt-Majumder et al. (2004), Wicks, Horowitz, Djordjevic & Gyulassy (2006), Peshier (2006): it is sizable! (either for heavy quarks only, for c only, for light quarks as well...)

Yet, at the same time...

Peigné, Gossiaux, Gousset (2005): yes, elastic energy loss is negligible, because the parton is formed inside the medium, not at infinity.

Conclusion... { ask Pol-Bernard and/or Thierry for further detail  
we'll know more at the time of the Deuxièmes Journées

