

Tutorial sheet 11

Discussion topic: What is a sound wave? How do you derive the corresponding equation of motion? How is the speed of sound defined? What happens when the amplitude of the wave becomes large?

21. One-dimensional “similarity flow”

Consider a perfect fluid at rest in the region $x \geq 0$ with pressure \mathcal{P}_0 and mass density ρ_0 ; the region $x < 0$ is empty ($\mathcal{P} = 0$, $\rho = 0$). At time $t = 0$, the wall separating both regions is removed, so that the fluid starts flowing into the region $x < 0$. The goal of this exercise is to solve this instance of *Riemann’s problem* by determining the flow velocity $\mathbf{v}(t, x)$ for $t > 0$. It will be assumed that the pressure and mass density of the fluid remain related by

$$\frac{\mathcal{P}}{\mathcal{P}_0} = \left(\frac{\rho}{\rho_0} \right)^\gamma, \quad \text{with } \gamma > 1$$

throughout the motion. This relation also gives you the speed of sound $c_s(\rho)$.

i. Assume that the dependence on t and x of the various fields involves only the combination $u \equiv x/t$.¹ Show that the continuity and Euler equations can be recast as

$$\begin{aligned} [u - \mathbf{v}(u)] \rho'(u) &= \rho(u) \mathbf{v}'(u) \\ \rho(u) [u - \mathbf{v}(u)] \mathbf{v}'(u) &= c_s^2(\rho(u)) \rho'(u), \end{aligned}$$

where ρ' resp. \mathbf{v}' denote the derivative of ρ resp. \mathbf{v} with respect to u .

ii. Show that the velocity is either constant, or obeys the equation $u - \mathbf{v}(u) = c_s(\rho(u))$, in which case the squared speed of sound takes the form $c_s^2(\rho) = c_s^2(\rho_0)(\rho/\rho_0)^{\gamma-1}$.

iii. Show that the results of **i.** and **ii.** lead to the relation

$$\mathbf{v}(u) = a + \frac{2}{\gamma - 1} c_s(\rho(u)),$$

where a denotes a constant whose value is fixed by the condition that $\mathbf{v}(u)$ remain continuous inside the fluid. Show eventually that in some interval for the values of u , the norm of \mathbf{v} is given by

$$|\mathbf{v}(u)| = \frac{2}{\gamma + 1} [c_s(\rho_0) - u].$$

iv. Sketch the profiles of the mass density $\rho(u)$ and the streamlines $x(t)$ and show that after the removal of the separation at $x = 0$ the information propagates with velocity $2c_s(\rho_0)/(\gamma - 1)$ towards the negative- x region, while it moves to the right with the speed of sound $c_s(\rho)$.

¹... which is what is meant by “self-similar”.