

Tutorial sheet 2

The exercise marked with a star is homework.

Discussion topics:

- What are the Lagrangian and Eulerian descriptions? How is a fluid defined?
- What are the strain rate tensor, the rotation rate tensor, and the vorticity vector? How do they come about and what do they measure?

*4. Stationary flow: second example

Consider the fluid flow whose velocity field $\vec{v}(t, \vec{r})$ has coordinates (in a given Cartesian system)

$$\mathbf{v}^1(t, \vec{r}) = kx^2, \quad \mathbf{v}^2(t, \vec{r}) = kx^1, \quad \mathbf{v}^3(t, \vec{r}) = 0, \quad (1)$$

where k is a positive real number, while x^1, x^2, x^3 are the coordinates of the position vector \vec{r} .

- i. Determine the stream lines at an arbitrary instant t .
- ii. Let X^1, X^2, X^3 denote the coordinates of some arbitrary point M and let t_0 be the real number defined by

$$kt_0 = \begin{cases} -\text{Artanh}(X^2/X^1) & \text{if } |X^1| > |X^2| \\ 0 & \text{if } X^1 = \pm X^2 \\ -\text{Artanh}(X^1/X^2) & \text{if } |X^1| < |X^2|. \end{cases}$$

Write down a parameterization $x^1(t), x^2(t), x^3(t)$, in terms of a parameter denoted by t , of the coordinates of *the* stream line $\vec{x}(t)$ going through M such that $d\vec{x}(t)/dt$ at any point *equals* the velocity field at that point, and that either $x^1(t) = 0$ or $x^2(t) = 0$ for $t = t_0$.

- iii. Viewing $\vec{x}(t)$ as the trajectory of a point—actually, of a fluid particle—, you already know the velocity of that point at time t (do you?). What is its acceleration $\vec{a}(t)$?
- iv. Coming back to the velocity field (1), compute first its partial derivative $\partial\vec{v}(t, \vec{r})/\partial t$, then the *material derivative*

$$\frac{D\vec{v}(t, \vec{r})}{Dt} \equiv \frac{\partial\vec{v}(t, \vec{r})}{\partial t} + [\vec{v}(t, \vec{r}) \cdot \vec{\nabla}]\vec{v}(t, \vec{r}).$$

Compare $\partial\vec{v}(t, \vec{r})/\partial t$ and $D\vec{v}(t, \vec{r})/Dt$ with the acceleration of a fluid particle found in question **iii**.

5. Yet another example of motion of a deformable continuous medium

Consider the motion defined in a system of Cartesian coordinates with basis vectors $(\vec{e}_1, \vec{e}_2, \vec{e}_3)$ by the velocity field with components

$$\mathbf{v}^1(t, \vec{r}) = f_1(t, x^2), \quad \mathbf{v}^2(t, \vec{r}) = f_2(t, x^1), \quad \mathbf{v}^3(t, \vec{r}) = 0,$$

with f_1, f_2 two continuously differentiable functions.

Compute the strain rate tensor $\mathbf{D}(t, \vec{r})$ for this motion. What is the volume expansion rate? Give the rotation rate tensor $\mathbf{R}(t, \vec{r})$ and the vorticity vector. Under which condition(s) on the functions f_1, f_2 does the motion become irrotational?

6. Isotropy of static pressure

Consider a geometrical point at position \vec{r} in a fluid at rest. The stress vector across every surface element going through this point is normal: $\vec{T}(\vec{r}) = -\mathcal{P}(\vec{r})\vec{e}_n$, with \vec{e}_n the unit vector orthogonal to the surface element under consideration. Show that the (hydrostatic) pressure \mathcal{P} is independent of the orientation of \vec{e}_n .

Hint: Consider the forces on the faces of an infinitesimal trirectangular tetrahedron.