

4. QCD in e^+e^- - annihilation

→ want to compare basic properties of (perturbative) QCD with experiment.

→ consider $e^+e^- \rightarrow$ hadrons

- total cross section $\sigma \sim \frac{\alpha_{em}^2}{s}$

calculate α_s corrections, $\sigma \rightarrow \sigma \cdot (1 + \alpha_s)$

renormalization scheme dependence enters at α_s^2

inclusive cross section: $(1 + \alpha_s + \alpha_s^2 + \alpha_s^3)$ known,
high precision QCD result!

non-perturbative corrections expected to be small

⇒ used as one of the most precise measurements of α_s .

- QCD predicts "jet" structure for final-state hadrons


define jet cross sections

calculate them, compare with experiment

⇒ can also be used to measure α_s ,
and to test/"see" triple-gluon-vertex.

4.1 $e^+e^- \rightarrow$ hadrons at leading order

reminder: 2→2 scattering in CMS (center of mass system)




detector measures in

$$d\Omega = \sin\theta \, d\theta \, d\phi$$

spherical coords.: $\int d\Omega = \int_0^\pi d\theta \sin\theta \int_0^{2\pi} d\phi = \int_{-1}^1 d(\cos\theta) \int_0^{2\pi} d\phi = 4\pi$

total cross section $\sigma = \int d\Omega \left(\frac{d\sigma}{d\Omega} \right) \leftarrow$ differential cross section

remember: Fermi's golden rule



$$\sigma_{2 \rightarrow n} = \frac{1}{F} \int d\Omega_n |M|^2$$

amplitude, e.g. from Feynman diagrams
phase space integral


$$\left(\prod_{i=1}^n \int \frac{d^3 p_i}{(2\pi)^3 2E_{p_i}} \right) (2\pi)^4 \delta^{(4)} \left(\sum_{i=1}^n p_i - q_A - q_B \right)$$

where

$$F = \frac{4 \sqrt{(q_A \cdot q_B)^2 - m_A^2 m_B^2}}{2 \sqrt{(s - m_A^2 - m_B^2)^2 - 4 m_A^2 m_B^2}} = 4(E_A + E_B) |\vec{q}_A|$$

$s = (q_A + q_B)^2 = (E_A + E_B)^2 - (\vec{q}_A + \vec{q}_B)^2$ in CMS, $q = (E, \vec{q})$

remember: amplitude $e^+ e^- \rightarrow \mu^+ \mu^-$ [see, e.g., Peskin/ Schroeder § 5.1]



$$= \bar{v}(q_B, s_B) (-ie\gamma^\mu) u(q_A, s_A) \left(\frac{-ig_{\mu\nu}}{q^2} \right) \bar{u}(p_1, s_1) (-ie\gamma^\nu) u(p_2, s_2)$$

\uparrow spin: +/- \uparrow Dirac spinor for incoming particle

start with unpolarized beam of e^+, e^-

\rightarrow average over spin states s_A, s_B : $\frac{1}{4} \sum_{s_A=\pm} \sum_{s_B=\pm}$

detector does not measure spin of final state

\rightarrow sum over spins s_1, s_2

$$\frac{1}{4} \sum_{s_i=\pm} |M|^2 = \frac{1}{4} \sum_s | \langle \text{final} | \text{initial} \rangle |^2 = \frac{1}{4} \sum_s \left| \frac{e^2}{q^2} \bar{v}_B \gamma^\mu u_A \bar{u}_1 \gamma^\nu v_2 \right|^2$$

$$= \frac{e^4}{4q^4} \sum_s \bar{v}_B \gamma^\mu u_A \bar{u}_1 \gamma^\nu v_2 \bar{u}_1 \gamma^\nu v_2 \bar{v}_B \gamma^\mu u_A$$

do spin sums via completeness rels

$$\sum_s u_A \bar{u}_A = \not{q}_A + m_e, \quad \sum_s v_2 \bar{v}_2 = \not{p}_2 - m_\mu$$

$$= \frac{e^4}{4q^4} \text{tr} \left((\not{q}_B - m_e) \gamma^\mu (\not{q}_A + m_e) \gamma^\nu \right) \text{tr} \left((\not{p}_1 + m_\mu) \gamma_\mu (\not{p}_2 - m_\mu) \gamma_\nu \right)$$


$$= \frac{e^4}{4q^4} 4 \left[q_B^\mu q_A^\nu + q_A^\mu q_B^\nu - g^{\mu\nu} (q_A \cdot q_B + m_e^2) \right] 4 \left[p_{1\mu} p_{2\nu} + p_{2\mu} p_{1\nu} - g_{\mu\nu} (p_1 \cdot p_2 + m_\mu^2) \right]$$

$$= \frac{8e^4}{q^4} \left(q_1 p_1 \cdot q_2 p_2 + q_1 p_2 \cdot q_2 p_1 + m_e^2 p_1 \cdot p_2 + m_\mu^2 q_A \cdot q_B + 2 m_e^2 m_\mu^2 \right)$$

CMS: $\vec{q}_B = -\vec{q}_A, \vec{p}_2 = -\vec{p}_1$; E-cons.: $2E = E_1 + E_2 = 2E_1$

$\Rightarrow E_A = E_B = E_1 = E_2 = E, \vec{q}_A^2 + m_e^2 = \vec{p}_1^2 + m_\mu^2$

$$= \frac{e^4}{E^4} \left(E^4 + m_e^2 E^2 + m_\mu^2 E^2 + (E^2 - m_e^2)(E^2 - m_\mu^2) \cos^2 \theta \right) \neq (\vec{q}_A, \vec{p}_1)$$


generalization: amplitude $e^+e^- \rightarrow f\bar{f}$ 

where $f \neq e$, $f \in \{u, d, s, c, b, \mu, \tau\}$

use charge $Q_f = \{0, 0, 0, +\frac{2}{3}, +\frac{2}{3}, +\frac{2}{3}, -1, -1, -\frac{1}{3}, -\frac{1}{3}, -\frac{1}{3}\}$

$$\Rightarrow \langle |M|^2 \rangle \equiv \sum_{\text{colors}} \sum_f \frac{1}{4} \sum_s |M|^2 = \sum_{\text{colors}} \frac{Q_f^2 e^4}{E^4} (E^4 + m_e^2 E^2 + m_f^2 E^2 + (E^2 - m_e^2)(E^2 - m_f^2) \cos^2 \theta)$$

reminder: phase space integration for 2→2 scattering



$$d\sigma_{2 \rightarrow 2} = \frac{1}{4} d\vec{E}_2 |M|^2, \quad \text{use } \delta^{(3)} \text{ for } \vec{p}_2 \text{-integral}$$

$$= \frac{1}{(8\pi)^2} \frac{|M|^2}{|\vec{q}_A| (E_A + E_B)} d^3_{\vec{p}_1} \frac{\delta(\sqrt{m_1^2 + \vec{p}_1^2} + \sqrt{m_2^2 + \vec{p}_1^2} - E_A - E_B)}{\sqrt{m_1^2 + \vec{p}_1^2} \sqrt{m_2^2 + \vec{p}_1^2}}$$

use that $|M|^2 = |M|^2(\vec{q}_A, \vec{q}_B, \vec{p}_1, \vec{p}_2) = |M|^2(|\vec{q}_A|, |\vec{p}_1|, \cos \theta)$

spherical coords, $s \equiv |\vec{p}_1|$, $d^3_{\vec{p}_1} = s^2 ds \sin \theta d\theta d\phi = d\Omega$

do $\int ds$ using δ fact

$$\frac{d\sigma_{2 \rightarrow 2}}{d\Omega} = \frac{1}{(8\pi)^2} \frac{|\vec{p}_1|}{|\vec{q}_A|} \frac{|M|^2(|\vec{q}_A|, |\vec{p}_1|, \cos \theta)}{(E_A + E_B)^2} \cdot \Theta(E - m_1 - m_2)$$

total cross section

$$\sigma = \int dR \frac{d\sigma}{dR} = \int_{-1}^1 d(\cos \theta) \frac{2\pi}{(8\pi)^2} \frac{\sqrt{E^2 - m_f^2}}{\sqrt{E^2 - m_e^2}} \frac{\langle |M|^2 \rangle}{(2E)^2} \Theta(E - m_f)$$

$$= \sum_{\text{colors}} \sum_f \frac{\pi}{3} \frac{Q_f^2 \alpha_{em}^2}{E^2} \frac{\sqrt{1 - m_f^2/E^2}}{\sqrt{1 - m_e^2/E^2}} \left(1 + \frac{m_e^2}{2E^2}\right) \left(1 + \frac{m_f^2}{2E^2}\right) \Theta(E - m_f)$$

$$\approx \frac{\pi}{3} \frac{\alpha_{em}^2}{E^2} \cdot \sum_{\text{colors}} \sum_f Q_f^2 \Theta(E - m_f), \quad \text{for } E \gg m_f \gg m_e$$

$$\Rightarrow R \equiv \frac{\sigma(e^+e^- \rightarrow \text{hadrons})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)} = \frac{\sum_f \sigma(e^+e^- \rightarrow q\bar{q})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)} \approx N_c \sum_f Q_f^2 \Theta(E - m_f)$$

$$\approx N_c \left\{ \left(\frac{2}{3}\right)_u^2 + \left(-\frac{1}{3}\right)_d^2 + \left(-\frac{1}{3}\right)_s^2 + \left(\frac{2}{3}\right)_c^2 + \left(-\frac{1}{3}\right)_b^2 + \left(\frac{2}{3}\right)_t^2 \right\} = N_c \left\{ \frac{4}{9}, \frac{5}{9}, \frac{2}{9}, \frac{10}{9}, \frac{11}{9}, \frac{1}{3} \right\}$$

